ROACH KILLER BAIT GEL

79432135 80215495F 200521AV1

FOR USE IN COMMERCIAL, INDUSTRIAL AND RESIDENTIAL AREAS

ACTIVE INGREDIENT: Fipronil*: [5-Amino-1-(2, 6-dichloro-4-trifluoromethyl) phenyl)-4-(1, R, S) -ftrifluoromethyl) sulfinyl)-1H-pyrazole-3-carbonitrilel 0.05% OTHER INGREDIENTS: 99.95% TOTAL:.... 100 000/ *CAS NO 120068-37-3

EPA Reg. No. 432-1460

EPA Est. No. 071106-GA-003

STOP - READ THE LABEL BEFORE USE. KEEP OUT OF REACH OF CHII DRFN CAUTION

See Panel for First Aid Instructions and Leaflet for Complete. Precautionary Statements and Directions for Use

Product may stain textiles and clothing.

FIDOT	

FIRST AID		
If on skin or clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.	
If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes	

 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment

advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin, Causes moderate eve irritation, Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PHYSICAL AND CHEMICAL HAZARDS: Do not use this product in or on electrical equipment where a possibility of shock hazard exists.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use MAXFORCE® FC MAGNUM ROACH KILLER BAIT GEL as a spot or crack and crevice treatment for the indoor and outdoor (adjacent to homes & structures) control of cockroaches in residential areas and the non-food/non-feed areas of institutional, warehousing and commercial establishments, warehouses, restaurants, food processing plants, supermarkets, hospitals, nursing homes, motels, hotels, apartment buildings, bakeries, beverage plants, breweries, bottling facilities, candy plants, canneries, cereal processing and manufacturing plants, dairies and dairy product processing plants, frozen food plants, schools, laboratories, computer facilities, sewers, "aircraft," buses, boats/ships, trains, homes, houses, industrial buildings, office buildings, kennels, kitchens, manufacturing facilities, spice plants, stores, wineries and similar structures, mausoleums, meat processing and packaging plants. meat and vegetable canneries, pet shops, and zoos. Non-food/non-feed areas include bedrooms, living rooms, bathrooms, closets, media rooms, garbage rooms, layatories, entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling). This product may also be used in food/feed areas of food/feed handling establishments as outlined below.

USE RESTRICTIONS

- Keep gel away from foods and food contact or preparation surfaces.
- Do not apply to areas where food/feed, utensils, or processing surfaces may become contaminated.
- USE ONLY IN AREAS NOT EASILY ACCESSIBLE TO CHILDREN, PETS, AND. OTHER NON-TARGET ORGANISMS.
- In pet shops and zoos place bait out of reach of pets and zoo animals. . Do not apply bait to surfaces where the temperature exceeds 130F as bait may run.
- Do not denosit del onto exposed surfaces.
- Do not apply bait gel to areas which have been recently sprayed with insecticide.
- . Do not spray insecticide over gel as it may cause the bait to become
- Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/prep and prepared food hold surfaces. Bait may be removed by washing.

USE DIRECTIONS

Place bait into cracks and crevices or openings leading into voids, or make spot applications near harborages, aggregation, foraging / feeding

Locations for bait placement; between different elements of construction; the hollow spaces in floors, walls, ceilings, or in equipment legs and bases; cracks and crevices around pipes, doors or windows, refrigerators, stoves, water heaters, dishwashers, and other equipment; behind or under baseboards, cupboards, cabinets, countertops, sinks; and, wherever cockroaches may be found hiding.

Numerous smaller placements will provide faster control than fewer larger spots, especially for German cockroach control.

Application rates for all cockroaches: Application rate is dependent on level of infestation and species to be controlled. Apply gel as round spots or thin beads into cracks and crevices or as a spot treatment.

• For use in maintenance programs where regular [monthly, quarterly]

- applications are made for continuous control of cockroach populations, use 0.75 grams of bait per square vard.
- . For light to moderate infestations (or for retreatments), use up to 1.5 grams of bait per square yard.
- To clean out severe or heavy infestations, use up to 3.0 grams of bait per square vard.

BAIT QUANTITY REFERENCE TABLE

Approximate weight per spot diameter	Approximate weight per bead length
0.1 gram = Pea size spot	1/3 gram = 1.5 inch bead
1.0 gram = Dime size spot	2/3 gram = 3.0 inch bead
1.5 gram = Quarter size spot	1 1/3 gram = 6 inch bead

APPLICATIONS IN FOOD/FEED HANDLING ESTABLISHMENTS

Application within food/feed areas of food/feed handling establishments is limited to crack and crevice treatment only. Do not deposit gel onto exposed surfaces; if gel contacts an exposed surface where food is handled, remove gel and wash exposed surface followed by a potable water rinse.

Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/prep and prepared food hold surfaces. Bait may be removed by washing.

Do not apply bait to surfaces where the temperature exceeds 130F as bait may run.

Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation. Application of this product in food/feed areas of food/feed handling establishments other than as a crack and crevice treatment is not permitted.

FOR OUTDOOR USE (adjacent to home or structure) Inject bait into cracks and crevices or expansion joints around windows and doors, porches, screens, eaves, sills, patios, garages, under stairways and other areas adjacent to structures where cockroaches may find harborage.

Refillable stations: MAXFORCE FC MAGNUM ROACH KILLER BAIT GEL may be placed into refillable bait stations designed for interior or exterior use; when placing balt in stations do not exceed 1 gm (175 teaspoon) and do not exceed the capacity of the stations. Close the station securely after application. During follow up visits inspect placements and reapply as needed

PANTRY PESTS: To control exposed, adult stages of beetles: cadelle, cigarette beetle, confused flour beetle, dermestid, drugstore beetle, flat grain beetle, granary weevil, lesser grain borer, lesser mealworm, mealworm, merchant grain beetle, red flour beetle, rice weevil, rusty grain beetle, saw-toothed grain beetle, spider beetle, or owarehouse beetle. After removing infested foodstuffs and cleaning up spilled foodstuffs and dust, applications of this product will kill adult stages of beetles remaining in area. Apply in voids, cracks and crevices, behind and under cupboards, and other difficult to reach areas where insects may hide. Application rate is deependent on level of infestation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE:** Store in a cool, dry area out of reach of

PESTICIDE DISPOSAL: If Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain

CONTAINER HANDLING: If Empty: Do not reuse container. Place in trash or offer for recycling if available.

If you have questions regarding the use of this product, please call 1-800-331-2867.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at

By using this product, user or buyer accepts the following Conditions. Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, plant injury, other property damage, as well as other unintended consequences may result because of factors beyond the control of Bayer CropScience LP Those factors include, but are not limited to, weather conditions, presence of other materials or the manner of use or application. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSOIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. NO agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE IP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd) and MAXFORCE® are registered trademarks of Bayer.



ROACH KILLER BAIT GEL

Note: Do not remove this label. Use only in areas not easily accessible to children and pets.

CAUTION:

Keep gel away from foods and food contact surfaces. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, diriking, chewing gum, or using tobacco.

Net Wt. 33g (1.16 Oz) 79432135 80211538F 200521AV1 Baver

CONTAINER HANDLING: If empty: Do not reuse container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. If you have questions regarding the use of this product, call toll free 1-800-331-29667.

May stain textiles, fabrics and/or porous surfaces. See Panel for First Aid Instructions and Leaflet for Complete Precautionary Statements and Directions for Use

Produced for:

Bayer Environmental Science A Division of Bayer CropScience LP 5000 CentreGreen Way, Suite 400 Cary, NC 27513 Product of China

Maxf file

BAYER R

FC MAGNUM ROACH KILLER BAIT GEL For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577 For PRODUCT USE Information Call 1-800-331-2867

Product may stain textiles and clothing.

 Take off contaminated clothing.
 Rinse skin immediately with plenty of water for 15-20 minutes.
 Call a poison control center or doctor for treatment advice. FIRST AID on skin r clothing:

 Hold eye open and rinse slowly and gently with water for 12-20 minutes. Permove contact lenses, if present, after the first 5 minutes, then continue rinsing, call a poison control center or doctor for treatment advice. •

in eyes:

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Hamful if absorbed through skin. Causes moderate eye irritation. Hamful if absorbed through skin, eyes, and cidhing, Wash throughly with soap and water after handling and before eating, drinking, chewing oum, or using tobacco. PHYSIGAL AND CHEMICAL HAZARDS: Do not use his product in every size, ALM CHEMICAL HAZARDS. Do not use this product in or on electrical equipment where a possibility of shock hazard exists

STORAGE AND DISPOSA.

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE. Store in a cool, dry area out of reach of children.

PESTICIDE DISPOSAL: if Partly Filled: Call your local solid waste agency for disposal instructions. Never place unused courtainter Hahbol. Mice. if Empty: Do not reuse container. Place in trast no cifer for recycling if available.

If you have questions regarding the use of this product, please call 1-800-331-2867.

See Panel for First Aid Instructions and Leaflet for Complete Precautionary Statements and Directions for Use.


Bayer

Produced for:
Bayer Environmental Science
A Division of Bayer CropScience LP
5000 CentreGreen Way, Suite 400
Cary, NO 27513
Product of China

MUNDAM 27 Maxfsrce









BAAB RERR

Open this end

ROACH KILLER BAIT GEL

Maxf
srce











ROACH KILLER BAIT GEL

.. 0.05% 99.95%

Total:*CAS NO. 120068-37-3

79432135 80183798F 200521AV1

EPA Reg. No. 432-1460 EPA Est. No. 071106-GA-003

STOP - READ THE LABEL BEFORE USE. KEEP OUT OF REACH OF CHILDREN. CAUTION

ROACH KILLER BAIT GEL

Contains:

1 Reservoir Net Weight 33g (1.16 Oz)



ROACH KILLER BAIT GEL